## Prime Farmland

(Only the soils considered prime farmland are listed. Urban or built-up areas of the soils listed are not considered prime farmland. If a soil is prime farmland only under certain conditions, the conditions are specified in parentheses after the soil name.)

Mitchell County, Iowa

```
Map
                                                    Soil name
symbol
27
       |Terril loam, 0 to 2 percent slopes
27B
       |Terril loam, 2 to 5 percent slopes
8.3
       |Kenyon loam, 0 to 2 percent slopes
83B
       |Kenyon loam, 2 to 5 percent slopes
       |Clyde silty clay loam, 0 to 3 percent slopes (Prime farmland if drained)
84
96
       |Turlin silt loam, 0 to 2 percent slopes (Prime farmland if protected from flooding or not
       | frequently flooded during the growing season)
98
       |Huntsville silt loam, 0 to 2 percent slopes
98B
       |Huntsville silt loam, 2 to 5 percent slopes
104B
       |Rockton silt loam, moderately deep, 2 to 5 percent slopes
       |Lamont fine sandy loam, 0 to 2 percent slopes
110
110B
       |Lamont fine sandy loam, 2 to 5 percent slopes
118
       |Garwin silty clay loam, 0 to 2 percent slopes (Prime farmland if drained)
119
       |Muscatine silty clay loam, 0 to 2 percent slopes
120
       |Tama silty clay loam, 0 to 2 percent slopes
120B
       |Tama silty clay loam, 2 to 5 percent slopes
135
       |Coland silty clay loam, 0 to 2 percent slopes (Prime farmland if drained and either protected
       | from flooding or not frequently flooded during the growing season)
151
       |Marshan clay loam, moderately deep, 0 to 2 percent slopes (Prime farmland if drained)
       |Marshan clay loam, deep, 0 to 2 percent slopes (Prime farmland if drained)
152
162B
       |Downs silt loam, 2 to 5 percent slopes
163B
       |Fayette silt loam, 2 to 5 percent slopes
       |Bassett loam, 0 to 2 percent slopes
171
171B
       |Bassett loam, 2 to 5 percent slopes
       |Dickinson fine sandy loam, 0 to 2 percent slopes
175
175B
       |Dickinson fine sandy loam, 2 to 5 percent slopes
177
       |Saude loam, 0 to 2 percent slopes
177B
       |Saude loam, 2 to 5 percent slopes
178
       |Waukee silt loam, 0 to 2 percent slopes
178B
       |Waukee silt loam, 2 to 5 percent slopes
183B
       |Dubuque silt loam, moderately deep, 2 to 5 percent slopes
184
       |Klinger silty clay loam, 0 to 2 percent slopes
       |Kensett silt loam, 0 to 2 percent slopes
188
198B
       |Floyd loam, 1 to 4 percent slopes
205B
       |Whalan loam, deep, 2 to 5 percent slopes
214B
       |Rockton loam, moderately deep, 2 to 5 percent slopes
       |Lawler loam, moderately deep, 0 to 2 percent slopes
225
226
       |Lawler loam, deep, 0 to 2 percent slopes
235
       |Coland-turlin complex, 0 to 2 percent slopes (Prime farmland if protected from flooding or not
       | frequently flooded during the growing season)
       |Bixby loam, 0 to 2 percent slopes
|Bixby loam, 2 to 5 percent slopes
265
2.65B
291
       |Atterberry silt loam, 0 to 2 percent slopes
302B
       \midCoggon loam, 2 to 5 percent slopes
303B
       |Pinicon loam, 1 to 4 percent slopes
       |Dinsdale silty clay loam, 0 to 2 percent slopes
377
377B
       |Dinsdale silty clay loam, 2 to 5 percent slopes
       |Maxfield silty clay loam, 0 to 2 percent slopes (Prime farmland if drained)
382
391B
       |Clyde-floyd complex, 1 to 4 percent slopes (Prime farmland if drained)
394
       |Ostrander loam, 0 to 2 percent slopes
394B
       |Ostrander loam, 2 to 5 percent slopes
398
       |Tripoli silty clay loam, 0 to 2 percent slopes (Prime farmland if drained)
399
       |Readlyn loam, 0 to 2 percent slopes
399B
       |Readlyn loam, 2 to 5 percent slopes
407B
       |Schley silt loam, 1 to 4 percent slopes (Prime farmland if drained)
```

USDA-NRCS, IOWA 04/03/2002

## Prime Farmland

(Only the soils considered prime farmland are listed. Urban or built-up areas of the soils listed are not considered prime farmland. If a soil is prime farmland only under certain conditions, the conditions are specified in parentheses after the soil name.)

## Mitchell County, Iowa

Map symbol	Soil name
482 482B 491B 507 536 575 575B 610B 714 714B 725 726	Oran loam, 0 to 2 percent slopes     Oran loam, 2 to 5 percent slopes     Racine silt loam, 0 to 2 percent slopes     Racine silt loam, 2 to 5 percent slopes     Racine silt loam, 2 to 5 percent slopes     Renova loam, 2 to 5 percent slopes     Canisteo silty clay loam, 0 to 2 percent slopes (Prime farmland if drained)     Hanlon sandy loam, 0 to 2 percent slopes     Dickinson-ostrander complex, 0 to 2 percent slopes     Dickinson-ostrander complex, 2 to 5 percent slopes     Lamont-renova complex, 2 to 5 percent slopes     Winneshiek loam, moderately deep, 0 to 2 percent slopes     Winneshiek loam, moderately deep, 2 to 5 percent slopes     Hayfield loam, moderately deep, 0 to 2 percent slopes     Hayfield loam, deep, 0 to 2 percent slopes     Calco silty clay loam, 0 to 2 percent slopes (Prime farmland if drained and either protected
761 771 771B 777 777B 781B 782 782B 783B 784B 797 798B 804 804B 805B 813 813B 904 904B	from flooding or not frequently flooded during the growing season)   Ansgar sitt loam, 0 to 2 percent slopes (Prime farmland if drained)   Franklin silt loam, 1 to 3 percent slopes   Waubeek silt loam, 0 to 2 percent slopes   Waubeek silt loam, 2 to 5 percent slopes   Wapsie loam, 0 to 2 percent slopes   Wapsie loam, 2 to 5 percent slopes   Sattre silt loam, 0 to 2 percent slopes   Soundes loam, 2 to 5 percent slopes   Donnan silt loam, 0 to 2 percent slopes   Donnan silt loam, 0 to 2 percent slopes   Cresco loam, 2 to 5 percent slopes   Gresco loam, 2 to 5 percent slopes   Grameston silty clay loam, 0 to 2 percent slopes   Jameston silty clay loam, 0 to 2 percent slopes   Ashdale silt loam, 0 to 2 percent slopes   Ashdale silt loam, 2 to 5 percent slopes   Roseville silt loam, 2 to 5 percent slopes   Roseville silt loam, 2 to 5 percent slopes   Askdalon loam, 0 to 2 percent slopes   Atkinson loam, 0 to 2 percent slopes   Atkinson loam, 0 to 2 percent slopes   Nasset silt loam, 0 to 5 percent slopes

USDA-NRCS, IOWA 04/03/2002